

ADAMYCIN LA 300

Oxytetracycline 300 mg/ml Injectable Solution For Animal Treatment only

Each ml contains:

Oxytetracycline Dihydrate B.P. equivalent to 300mg of Oxytetracycline base in an aqueous solution.

Dosage And Administration:

Adamycin LA 300 can only be administered as a single dose either at a rate of 20mg/kg for 3 to 4 days duration of activity or 30mg/kg for prolonged duration of activity (i.e. activity maintained for 5 to 6 days).

Cattle and Pigs:

Standard dose: 20mg/kg (1 ml/ 15kg) High dose - 30 mg/kg (1ml/10kg)

Administer by deep intramuscular injection to cattle and pigs for the treatment and control of conditions caused by or associated with organisms sensitive to Oxytetracycline.

Withholding times.

Meat: Animals must not be slaughtered for human consumption during treatment

20 mg/kg dose: Cattle may be slaughtered for human consumption only after 28 days from the last treatment.

Pigs may be slaughtered for human consumption only after 14 days from the last treatment.

30 mg/ kg dose: Cattle may be slaughtered for human consumption only after 35 days from the last treatment.

Pigs may be slaughtered for human consumption only after 28 days from the last treatment.

Milk: Milk for human consumption must not be taken during treatment. Milk for human consumption may be taken only from cows after 7 days from the last treatment.

Following withdrawal of the first dose use the product within 28 days.

Discard unused material

Keep out of reach of children.

Use a sterile needle and syringe.

STORE AT A ROOM TEMPERATURE. (not exceeding 25°C).

FOR ANIMAL TREATMENT ONLY.

Protect from light.

Manufactured by:

Norbrook Laboratories Ltd

Station Works, Newry Co. Down

Northern Ireland, U.K.

POM



BIMEDA LTD.

Funzi Road, Industrial Area. P.O. Box 30620-00100,

Nairobi, Kenya

Tel: +254 20 6537622-6 Fax: +254 20 6537628