

# BUVONAL injection

For treatment of Theileriosis in cattle, including infection with East Coast Fever.

## COMPOSITION:

Buvonal contains Buparvaquone 50 mg per ml. Buparvaquone is an analogue of Parvaquone, a hydroxynaphthoquinone. It has a greater resistance to metabolic degradation than Parvaquone, resulting in much higher activity than Parvaquone.

## INDICATIONS

Buvonal injection is indicated for the treatment of Theileriosis in cattle. It is highly effective against Theileria species including the most significant Theilerial pathogens *T. parva* (*parva*), *T. parva* (*lawrencei*) *T. parva* (*bovis*), *T. annulata*, *T. mutans* and *T. orientalis* (*sergenti*).

## DOSAGE AND ADMINISTRATION:

Intramuscular injection into the neck muscles.

Calves and adult cattle: 1 ml per 20 kg bodyweight (2.5 mg/kg)

Bodyweight	Dose
100 kg	5 ml
150 kg	7.5 ml
200 kg	10 ml
250 kg	12.5 ml
300 kg	15 ml

More advanced cases may require 1 or more additional injections given at an interval of 2 to 3 days as clinically indicated.

## WITHDRAWAL PERIOD:

Meat: 42 days.

Milk: 48 hours.

## Storage:

Store between 2°C and 25°C.

Protect from light.

Keep out of reach of children.

