

# CURAFLUKE

**A 5% w/v oral suspension of Fenbendazole**

**Ph. Eur 50 mg/ml and Rafoxanide B.P. (Vet) 50mg/ml.**

**Curaf Luke** is active against Fluke, Lungworms, Tapeworms, Stomachworms in cattle and sheep plus Nasal bots (*Oestrus ovis*) in sheep. It is a broad spectrum anthelmintic for the treatment of *Oestrus ovis*, *Haemonchus sp.*, *Ostertagia sp.*, *Trichostrongylus sp.*, *Cooperia sp.*, *Nematodirus sp.*, *Bunostomum sp.*, *Trichuris sp.*, *Strongyloides sp.*, *Dictyocaulus sp.*, *Moniezia sp.*, *Fasciola sp.* Curaf Luke has a good therapeutic effect against type II *Ostertagiasis*.

To be given orally at a dose rate of 7.5 mg/kg bodyweight for sheep and 11.25 mg/kg bodyweight for cattle.



## DOSAGE & ADMINISTRATION

The dose for heavier animals is an additional 11.25 ml per 50 kg for cattle and 1.5 ml per 10 k.g for sheep

Cattle	Dose ml	Dose/Pack	Sheep	Dose ml	Dose/Pack
50 kg	12 ml	83	10 kg	1.50 ml	666
100 kg	23 ml	43	15 kg	2.25 ml	444
150 kg	34 ml	29	20 kg	3.00 ml	333
200 kg	45 ml	22	25 kg	3.75 ml	266
250 kg	57 ml	17	30 kg	4.50 ml	222

## WITHHOLDING PERIOD

Not for use in dairy cows during lactation. Animals must not be slaughtered for human consumption during treatment or for 28 days thereafter. Store in a cool place. Avoid the introduction of contamination during use. **For animal Treatment Only.**

**Available Packs : 60ml, 120ml, 250ml, 500ml, 1 Litre**

**BIMEDA LTD.**

Funzi Road, Industrial Area. P.O. Box 30620-00100,  
Nairobi, Kenya  
Tel: +254 20 6537622-6 Fax: +254 20 6537628

[www.bimeda.com](http://www.bimeda.com)