

MULTIMAST

DRY COW INTRAMAMMARY SUSPENSION

Neomycin Sulphate 100mg / Penethamate Hydriodide 100mg / Procaine Benzylpenicillin 400mg

DATA SHEET



INDICATIONS

Multimast Dry Cow intramammary suspension is licensed for routine use in cows at drying off, to treat existing intramammary infections and to assist in preventing new infections occurring during the dry period.

MORE

Each syringe (4.5g intramammary suspension) contains:

- Neomycin Sulphate 100mg
- Penethamate Hydriodide 100mg
- Procaine Benzylpenicillin 400mg



LIST No	UNIT PACKAGE	CASE SIZE
1MUL004	120 Syringes	1



See reverse for Administration & Dosage

Multimast

DRY COW INTRAMAMMARY SUSPENSION



Neomycin Sulphate 100mg
Penethamate Hydriodide 100mg
Procaine Benzylpenicillin 400mg

PRESENTATION

Single dose plastic syringe containing a smooth off-white oily suspension for intramammary administration to cows.

Each 4.5g intramammary suspension contains:

Neomycin Sulphate 100mg
Penethamate Hydriodide 100mg
Procaine Benzylpenicillin 400mg

INDICATIONS

For routine use in cows at drying off, to treat existing intramammary infections and to assist in preventing new infections occurring during the dry period.

DOSAGE & ADMINISTRATION

The contents of one syringe should be infused into each quarter via the teat canal immediately after the final milking of a lactation. Before infusion, the teat should be thoroughly cleaned and disinfected and care should be taken to avoid contamination of the injector nozzle. Following infusion, it is advisable to use a teat dip or spray.

CONTRA-INDICATIONS, WARNINGS, ETC

For animal treatment only.
Do not use in the lactating cow.
Do not use within 50 days prior to calving.
Do not use in animals with known hypersensitivity to the active ingredients.

OPERATOR WARNINGS

Penicillin and cephalosporins may cause sensitisation following injection, inhalation, ingestion or skin contact. Sensitivity to penicillin may lead to cross reactions to cephalosporins and vice versa. Allergic reactions to these substances may occasionally be serious.

1- Do not handle this product if you know that you are sensitised, or if you have been advised not to work with such preparations.

2- Handle this product with care to avoid exposure, taking all recommended precautions.

3- If you develop symptoms such as skin rash following exposure, seek medical advice and show this warning to the doctor. Swelling of the face, lips or eyes, or difficulty with breathing are more serious symptoms and require urgent medical attention.

4- Any unused product or waste material should be disposed of in accordance with current practice for pharmaceutical waste under national waste disposal regulations.

WITHDRAWAL PERIODS

Milk for human consumption may only be taken after 96 hours post calving in cows with a dry period of more than 50 days.

Milk for human consumption may only be taken after 50 days plus 96 hours after the last treatment for cows with a dry period of 50 days or less.

Animals may not be slaughtered for human consumption until 28 days from the treatment.

PHARMACEUTICAL PRECAUTIONS

Keep out of reach and sight of children.
Do not store above 25°C.

SHELF LIFE

Shelf life 2 years
Single use syringes. Discard after use.

LEGAL CATEGORY

POM

PACKAGE QUANTITIES

120 x 4.5g injectors

MARKETING AUTHORISATION NUMBER

VPA 22033/030/001

MARKETING AUTHORISATION HOLDER

Bimeda® Animal Health Limited
2, 3 & 4 Airton Close, Airton Road
Tallaght, Dublin 24, Ireland
Tel: +353 (01) 466 7900

A full product SPC is available on request from Bimeda or alternatively can be found on the HPRA website.

Use Medicines Responsibly.

TAKE TIME



OBSERVE LABEL
DIRECTIONS

www.bimeda.ie