

Global Excellence in Animal Health

NITRODOX

34% Nitroxynil Solution for injection

Presentation:

Nitrodox is presented for use in cattle, sheep and goats as a solution containing 34% w/v Nitroxynil.

Indications

Nitrodox is indicated in the treatment of Mature and Immature Liver Fluke infestation, Parafilaria bovicola (false bruising) and Roundworms in cattle; and wire worm, nodular worm, nasal worm and Liver Fluke in sheep and goats.

1. Cattle:

Gastrointestinal nematodes

- Wire worm
- Hookworm
- Nodular worm

Trematodes

Liver fluke (F. hepatica & F. gigantica)

2. Sheep and goats

Gastrointestinal nematodes

- Wireworm
- Nodular worm

Trematodes

- Liver fluke (F. hepatica)
- Nasal worm (1st, 2nd &3rd instar stages)

Dosage and Administration

Use only as directed by subcutaneous injection.

1, Liverfluke and roundworms other than Parafilaria: standard dose.1.5 ml per 50 kg body weight (10mg/kg)

2. Parafilaria bovicola (False bruising): double dose. 3 ml per 50 kg body (20 mg/kg)

Live	Liverfluke and roundworms	Parafilaria
weight	Standard dose 10mg/kg	Double dose 20 mg/kg
50 kg	1.5 ml	3.0 ml
100 kg	3.0 ml	6.0 ml
150 kg	4.5 ml	9.0 ml
200 kg	6.0 ml	12.0 ml

If total dose exceeds 10 ml, then inject in more than one place and massage injection site thoroughly.



SHEEP AND GOATS

Use only as directed by subcutaneous injection. Liver fluke, wireworm, nodular worm and nasal worm.

1 ml per 25 kg body weight (13.6 mg/kg)

Live weight	Standard dose
15 kg	0.6 ml
20 kg	0.8 ml
25 kg	1 ml
50 kg	2 ml
60 kg	2.4 ml

WITHDRAWAL PERIOD

MEAT:

- Animals treated at standard dose: 60 days for cattle and Goats, 45 days for sheep.
- Animals treated for Parafilaria at double dose: 70 days.

MILK:

- Animals treated at standard dose: 3 days.
- Do not treat lactating cows at double dose if the milk is used for human consumption.

PRECAUTIONS

- Do not exceed the recommended dosage.
- Do not treat calves younger than 6 months with the double dose.
- Keep out of reach of children, un-informed persons and animals
- For animal treatment only.
- Store below 25°c.
- Protect from light

Manufactured by:

Chongqing Fangtong Animal Pharma. Co. Ltd. Chongqing, China.

Under license from:

BIMEDA Broomhill Road, Tallaght, Dublin 24, Ireland.