

Tramazole 10%

A 10% suspension of Albendazole 100 mg, Methylparaben 2 mg & Propylparaben 0.2 mg

Protects against round, Tape and fluke worms

The recommended use Tramazole 10% is a broad spectrum multi-purpose anthelmintic for control of mature and developing immature forms of gastrointestinal roundworms, lungworms, tapeworms and adult liver fluke in sheep and cattle. The product is also ovicidal against fluke and roundworm eggs.

In sheep it is active against benzimidazole-susceptible strains of the following species:

- Roundworms:** Ostertagia, Haemonchus, Trichostrongylus, Nematodirus (including N. battus), Chabertia and Oesophagostomum.
It is usually effective against inhibited larvae of Ostertagia.
- Lungworms:** Dictyocaulus filaria.
- Tapeworms:** Moniezia spp.
- Adult Liver Fluke:** Fasciola hepatica.

In cattle it is active against the following species:

- Roundworms:** Ostertagia, Haemonchus, Bunostomum, Trichostrongylus, Nematodirus, Oesophagostomum, Bunostomum, Cooperia and Strongyloides spp.
It is usually effective against inhibited larvae of Cooperia and Ostertagia.
- Lungworms:** Dictyocaulus viviparus.
- Tapeworms:** Moniezia spp.
- Adult Liver Fluke:** Fasciola hepatica.

Dosage and Administration:

For oral administration only using properly calibrated dosing equipment. Estimate bodyweight accurately.

Sheep: For the control of roundworms, lungworms, tapeworms and fluke and roundworm eggs, 5 mg Albendazole per kg bodyweight which is equivalent to 1.0 ml Tramazole 10% per 20 kg bodyweight. For the additional treatment of adult liver fluke (chronic fascioliasis) in sheep, 7.5 mg Albendazole per kg bodyweight which is equivalent to 1.5 ml Tramazole 10% per 20 kg bodyweight.

Cattle: For the control of roundworms, lungworms, tapeworms and fluke and roundworm eggs, 7.5 mg Albendazole per kg bodyweight which is equivalent to 7.5 ml Tramazole 10% per 100 kg bodyweight. For the additional treatment of adult liver fluke (chronic fascioliasis) in cattle, 10 mg Albendazole per kg bodyweight which is equivalent to 10 ml Tramazole 10% per 100 kg body weight.

DOSAGE GUIDE

DOSAGE	ROUND AND TAP EWORMS	ADULT LIVER FLUKE
SHEEP/ GOATS	2.5ml/50kg Bodyweight or 5.0mg/kg Bodyweight.	3.7ml/50kg Bodyweight or 7.5mg/kg Bodyweight.
CATTLE/ CAMEL	3.7ml/50 kg Bodyweight or 7.5mg/100 kg Bodyweight. 7.5mg/kg Bodyweight.	5.0ml/50kg Bodyweight or 10ml/100kg Bodyweight or

DO NOT ADMINISTER DURING FIRST 3 WEEKS OF PREGNANCY



Available Packs : 40ml, 120ml, 500ml and 1Lt

BIMEDA LTD.

Funzi Road, Industrial Area. P.O. Box 30620-00100,
Nairobi, Kenya
Tel: +254 20 6537622-6 Fax: +254 20 6537628