

Parvexon Plus Inj. (Parvaquone + Frusemide)

**PARVEXON PLUS™
INJECTION**

FOR ECF WITH PULMONARY OEDEMA (PARVAQUONE + FRUSEMIDE)

DESCRIPTION: A potent injection treatment for theileriosis (East Coast Fever, ECF), particularly for advanced cases with pulmonary oedema; containing in each ml, parvaquone 150mg and frusemide 55mg. Parvaquone is theilericidal, acting on schizonts and piroplasms; Frusemide is a diuretic, which resolves pulmonary oedema.

DOSAGE AND ADMINISTRATION: Inject by the intramuscular route into the neck muscles at the rate of 1ml/15kg (10.0 mg Parvaquone, 3.6 mg Frusemide per kg). Repeat after 48 hours. In cases of exceptionally severe infections with pulmonary oedema, further treatment at half the dosage rate may be required at 24 hour intervals.

Calves: 1ml per 15kg bodyweight

Young-stock and adults:

Bodyweight	Dose
100kg	7 ml
150kg	10 ml
200kg	14 ml
250kg	17 ml
300kg	20 ml

Alternatively, when pulmonary symptoms have been resolved by Parvexon Plus, treatment to achieve parasitic cure may be completed with Parvexon, which contains only parvaquone, at a dose rate of 1ml/15kg every 48 hours.

Do not use by the intravenous or subcutaneous routes. Normal aseptic precautions should be observed.

CONTRA-INDICATIONS AND WARNINGS: East Coast Fever (ECF) is an immuno-depressive disease and potentially fatal concurrent infections are common. In particular, anaplasmosis may be activated from the carrier state and the principal lesions of ECF, such as oedema of the lungs and the alimentary canal, predispose to bacterial pneumonia and enteritis. The symptoms of these conditions may be masked by those of ECF. Special care should be taken to diagnose any complicating conditions and to treat them promptly and specifically. Adverse reaction to parvaquone is extremely rare. Frusemide is chemically similar to the sulphonamides so Parvexon Plus should not be used in animals with sulphonamide sensitivity. Local swelling may occur at the injection site but this is usually transient.

**Milk withholding period – 14 days
Meat withholding period – 28 days**

**FOR ANIMAL TREATMENT ONLY.
KEEP OUT OF REACH OF CHILDREN.
STORE BELOW 25°C.
PROTECT FROM LIGHT**

50ml glass vial, 100ml glass vial



BIMEDA LTD.

Funzi Road, Industrial Area. P.O. Box 30620-00100,
Nairobi, Kenya
Tel: +254 20 6537622-6 Fax: +254 20 6537628